100 deg^2 Mock Galaxy Cone for H I Surveys with the Early SKA

D. Obreschkow and M. Meyer

International Centre for Radio Astronomy Research (ICRAR)

June 5, 2014

arXiv:1406.0966v1 [astro-ph.GA] 4 Jun 2014

This document accompanies an easy-to-use mock catalog of galaxies with detailed neutral *atomic* hydrogen (H I) and auxiliary *molecular* and *optical* properties. The catalog covers a field of 10-by-10 degrees and a redshift range of $z = 0-1.2$. It contains galaxies with 21 cm peak flux densities down to 1 μJy and is, within this flux limit, complete for H I masses above $10^8 M_\odot$. Five random realisations of the catalog in ASCII format (∼4 GB/file) and subtables with H I flux limits of 10 μJy (∼500 MB/file) and 100 μJy (∼30 MB/file) can be downloaded at

http://ict.icrar.org/store/staff/do/s3sax

Context: This catalog was released in the frame of the 2014 renewal of the science case for the Square Kilometre Array (SKA) telescope. It is the basis for performance calculations in several chapters of this science case and can be considered a reference tool to assess future surveys with the SKA and its pathfinders.

Model: The catalog is a light version of the Semi-Analytic Suite of the SKA Simulated Skies (S^3-SAX), delivered as part of the European SKA design studies (SKADS) in 2009. It relies on the physical models described in Obreschkow et al. [2009a,b,c], applied to the *evolving* semi-analytic model by De Lucia & Blaizot [2007] on the Millennium dark matter simulation [Springel et al., 2005]. More sophisticated models and models run with updated cosmological parameters have since been produced, but the differences to S^3-SAX are minor in the considered redshift range. The S^3-SAX model has the advantage of having been exhaustively tested in more than a hundred peer-reviewed publications, successfully reproducing H I masses, line profiles, Tully-Fisher relation, disk sizes, angular momentum, clustering and evolution characteristics.

Cosmic volume: The volume of the catalog is a cone with a 10-by-10 degree opening, truncated at a redshift of $z = 1.2$. For H I, this z limit matches the 650 MHz lower limit of SKA1-Survey band 2 specified in the System Baseline Design (March 2013). Five independent cones are available to assess the cosmic variance within this volume. They are centred on the coordinates RA = 36/108/180/252/324° and Dec = −30°.

Flux/mass limit: The catalog has been truncated to 21 cm peak flux densities above 1 μJy to keep the catalog size reasonable. Within this flux limit, the model is complete for H I masses above $10^8 M_\odot$. At smaller H I masses, the number of galaxies drops steeply, missing most of the H I in dwarf galaxies. This limit is only a minor concern when dealing with isolated direct H I detections in blind surveys at $z > 0$, where only a tiny fraction of the total survey volume is sensitive to H I masses $< 10^8 M_\odot$. However, when dealing with global H I mass estimates, e.g. in a stacking experiment to measure $\Omega_{\rm H\,I}(z)$, the H I mass contained in unresolved galaxies is almost certainly significant (cf. Section 3.4 of Obreschkow et al., 2011), as apparent from direct comparisons of the modelled $\Omega_{\rm H\,I}(z > 1)$ against inferences from DLA data, hydrodynamical simulations by Pontzen et al. [2008] and statistical extrapolations by Lagos et al. [2011]. It is recommended that applications relying on the *total* H I content of the universe at $z \sim 1$ correct for the H I mass resolution limit of the catalog – most simplistically by scaling all H I masses in the mock catalog with a z-dependent factor (> 1) to match $\Omega_{\rm H\,I}(z)$ inferred from the best DLA and/or stacking data.

Optical properties: Whereas the original S^3-SAX release did not contain optical properties, this new catalog includes approximate optical radii, stellar masses and apparent Vega R-band magnitudes, derived from simulated extinction corrected absolute magnitudes using a variable K-correction, fitted to empirically determined K-corrections as a function of redshift and morphology [Driver et al., 1994, Westra et al., 2010]. In a test sample of 10^3 mock galaxies, these apparent magnitudes were consistent with the 'exact' magnitudes obtained by redshifting empirical SEDs fitted to the simulated absolute Vega BVRIK magnitudes.

Need additional properties? This catalog has intentionally been limited to 20 properties per galaxy. Additional properties listed on http://s-cubed.physics.ox.ac.uk/s3_sax/sky (e.g. alternative CO line properties, surface density profiles, metallicity, black hole properties) can be downloaded there, respectively requested directly from the authors of this document.

We acknowledge Simon Driver for his hands-on assistance in calculating the K-corrections.

Col	Symbol	Unit	Description
1	ID	–	Unique galaxy identifier in the Munich Semi-Analytic Model "DeLucia2006a"
2	RA	deg	Right ascension of galaxy centre
3	Dec	deg	Declination of galaxy centre
4	z	–	Apparent redshift of galaxy centre, including the Doppler component due to peculiar motion relative to the Hubble expansion
5	i	deg	Galaxy inclination defined as the smaller angle ($0° - 90°$) between the line-of-sight and the rotational axis of the galaxy
6	T	–	Numerical Hubble type (-6...0 for ellipticals, 0...10 for spirals, 99 for morphologically unresolved objects, mostly dwarfs)
7	M_*	$\mathrm{M}_\odot$	Stellar mass
8	$M_{\mathrm{H\,I}}$	$\mathrm{M}_\odot$	Mass of neutral atomic hydrogen H I, without helium
9	M_{H_2}	$\mathrm{M}_\odot$	Mass of molecular hydrogen H_2, without helium
10	$S_{\mathrm{H\,I}}^{\mathrm{int}}$	$\mathrm{Jy\,km\,s^{-1}}$	Velocity-integrated flux of the redshifted 21 cm H I emission line, with velocity units defined in the galaxy rest-frame
11	$S_{\mathrm{H\,I}}^{\mathrm{peak}}$	Jy	Peak flux density of the H I emission line; typically the flux density of the 'horns'
12	$S_{\mathrm{CO}}^{\mathrm{int}}$	$\mathrm{Jy\,km\,s^{-1}}$	Velocity-integrated flux of the redshifted 115.27 GHz ^{12}CO(1–0) emission line, with velocity units defined in the galaxy rest-frame
13	$S_{\mathrm{CO}}^{\mathrm{peak}}$	Jy	Peak flux density of the ^{12}CO(1–0) emission line; typically the flux density of the 'horns'
14	$W_{\mathrm{H\,I}}^{50}$	$\mathrm{km\,s^{-1}}$	Width of the H I emission line, in galaxy rest-frame velocity units, measured at 50% of the peak flux density
15	$W_{\mathrm{H\,I}}^{20}$	$\mathrm{km\,s^{-1}}$	Width of the H I emission line, in galaxy rest-frame velocity units, measured at 20% of the peak flux density
16	$r_{\mathrm{H\,I}}^{\mathrm{edge}}$	arcsec	Apparent H I radius along the major axis out to a H I disk surface density of $1\,\mathrm{M}_\odot\mathrm{pc}^{-2}$, corresponding to a face-on column density of $1.25 \cdot 10^{20}\mathrm{cm}^{-2}$
17	$r_{\mathrm{H\,I}}^{\mathrm{half}}$	arcsec	Apparent H I half-mass radius along the major axis
18	M_{R}	mag	Absolute Vega R-band magnitude, corrected for intrinsic dust extinction; 99 if stellar mass and star formation history are insufficiently resolved to compute M_{R}
19	m_{R}	mag	Apparent Vega R-band magnitude; value 99 if no absolute magnitudes available
20	r_{e}	arcsec	Effective radius, here approximated as the radius containing half the stellar mass if the galaxy were viewed face-on

Table 1: Description of the columns of mock catalog in ASCII format.

References

De Lucia G., Blaizot J., 2007, MNRAS, 375, 2

Driver S. P., Phillipps S., Davies J. I., Morgan I., Disney M. J., 1994, MNRAS, 266, 155

Lagos C., et al., 2011, MNRAS, submitted

Obreschkow D., Croton D., DeLucia G., Khochfar S., Rawlings S., 2009a, ApJ, 698, 1467

Obreschkow D., Heywood I., Klöckner H.-R., Rawlings S., 2009b, ApJ, 702, 1321

Obreschkow D., Heywood I., Rawlings S., 2011, ApJ, 743, 84

Obreschkow D., Klöckner H., Heywood I., Levrier F., Rawlings S., 2009c, ApJ, 703, 1890

Pontzen A., et al., 2008, MNRAS, 390, 1349

Springel V., et al., 2005, Nat, 435, 629

Westra E., Geller M. J., Kurtz M. J., Fabricant D. G., Dell'Antonio I., 2010, PSPS, 122, 1258



Figure 1: Representation of the five statistically independent mock cones of galaxies. The cones are centred on the coordinates RA = 36/108/180/252/324° and Dec = −30°.



Figure 2: Redshift distribution of galaxies, where the y-axis shows dN/dz per *square degree*, averaged from the 100 square degree field of view. The five colours represent the five cones shown in Fig. 1. LEFT: all galaxies in the full ($S_{\rm H\,I}^{\rm peak} \geq 1\,\mu$Jy) cones. RIGHT: all galaxies with $S_{\rm H\,I}^{\rm peak} \geq 100\,\mu$Jy (thin lines) and subsamples of these galaxies (thick lines) with an apparent R-band magnitude $m_{\rm R} \leq 20$ and an effective radius $r_{\rm e} \geq 1''$, as an example of matching H I data to a GAMA-like optical survey to derive approximate stellar mass and inclination measurements.